

CARB-COUNTING? GLUTEN-FREE?

Yet still not feeling great?


When you're diagnosed with both Celiac Disease and Type 1 Diabetes, there is so much to handle. Eating can turn into a challenging, misunderstood, and lonely struggle.

Don't go it alone anymore.

Be seen by specialists when you join **T1D+CeD** - a research study for those diagnosed with both Type 1 Diabetes and Celiac Disease. Try a new investigational medication designed to reduce symptoms of accidental gluten exposure.

Study Details

- 6 visits over 22 weeks
- Drink-based study medication
- CGM provided for 2 weeks
- HbA1c taken at most visits
- No biopsies or injections involved
- Weekly snacks provided
- Payment for participation & travel



Qualifications:

- Diagnosed with both Type 1 Diabetes and Celiac Disease
- Experienced moderate abdominal pain, tiredness, and/or bloating symptoms in the past 30 days
- Between 18-80 years old
- On a gluten-free diet for ≥ 12 months
- No history of IBS, IBD, or gastroparesis

T1D  **CeD**TM

Diabetes+Celiac Research Study

For participant's rights questions,
contact 1-866-680-2906.

Scan to
pre-screen now:



For more information, contact:

Nora Arrizon-Ruiz
Stanford University

narrizon@stanford.edu

650-721-8624